



CTDP1 gene

CTD phosphatase subunit 1

Normal Function

The *CTDP1* gene provides instructions for making a protein called carboxy-terminal domain phosphatase 1. This protein helps regulate the activity of an enzyme called RNA polymerase II. The RNA polymerase II enzyme initiates transcription, which is a key step in using the information carried by genes to direct the production (synthesis) of proteins.

Health Conditions Related to Genetic Changes


congenital cataracts, facial dysmorphism, and neuropathy

Thus far, everyone with CCFDN has had the same mutation in both copies of the *CTDP1* gene in each cell. This mutation, written as IVS6+389C>T, alters the way the gene's instructions are pieced together to produce the carboxy-terminal domain phosphatase 1 protein. The altered instructions introduce a premature stop signal, resulting in an abnormally short, nonfunctional protein that cannot regulate transcription. Defective regulation of the transcription process affects the development and function of many parts of the body. It is not known how nonfunctional carboxy-terminal domain phosphatase 1 protein results in the specific signs and symptoms of CCFDN.

Chromosomal Location

Cytogenetic Location: 18q23, which is the long (q) arm of chromosome 18 at position 23

Molecular Location: base pairs 79,679,801 to 79,754,510 on chromosome 18 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- CCFDN
- CTD (carboxy-terminal domain, RNA polymerase II, polypeptide A) phosphatase, subunit 1
- CTD of POLR2A, phosphatase of, subunit 1
- CTDP1_HUMAN
- FCP1
- RNA polymerase II subunit A C-terminal domain phosphatase
- RNA polymerase II subunit A C-terminal domain phosphatase isoform FCP1a
- RNA polymerase II subunit A C-terminal domain phosphatase isoform FCP1b
- serine phosphatase FCP1a
- TFIIF-associating CTD phosphatase 1
- transcription factor IIF-associating CTD phosphatase 1

Additional Information & Resources

Educational Resources

- Molecular Biology of the Cell (fourth edition, 2002): RNA Polymerase II Requires General Transcription Factors
<https://www.ncbi.nlm.nih.gov/books/NBK26887/#A999>

GeneReviews

- Congenital Cataracts, Facial Dysmorphism, and Neuropathy
<https://www.ncbi.nlm.nih.gov/books/NBK25565>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28CTDP1%5BTIAB%5D%29+OR+%28%28FCP1%5BTIAB%5D%29+OR+%28CCFDN%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+2520+days%22%5Bdp%5D>

OMIM

- C-TERMINAL DOMAIN OF RNA POLYMERASE II SUBUNIT A, PHOSPHATASE OF, SUBUNIT 1
<http://omim.org/entry/604927>

Research Resources

- **Atlas of Genetics and Cytogenetics in Oncology and Haematology**
http://atlasgeneticsoncology.org/Genes/GC_CTDP1.html
- **ClinVar**
<https://www.ncbi.nlm.nih.gov/clinvar?term=CTDP1%5Bgene%5D>
- **HGNC Gene Family: CTD family phosphatases**
<http://www.genenames.org/cgi-bin/genefamilies/set/1041>
- **HGNC Gene Symbol Report**
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=2498
- **NCBI Gene**
<https://www.ncbi.nlm.nih.gov/gene/9150>
- **UniProt**
<http://www.uniprot.org/uniprot/Q9Y5B0>

Sources for This Summary

- OMIM: C-TERMINAL DOMAIN OF RNA POLYMERASE II SUBUNIT A, PHOSPHATASE OF, SUBUNIT 1
<http://omim.org/entry/604927>
- GeneReview: Congenital Cataracts, Facial Dysmorphism, and Neuropathy
<https://www.ncbi.nlm.nih.gov/books/NBK25565>
- Kalaydjieva L. Congenital cataracts-facial dysmorphism-neuropathy. *Orphanet J Rare Dis.* 2006 Aug 29;1:32. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16939648>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1563997/>
- Majello B, Napolitano G. Control of RNA polymerase II activity by dedicated CTD kinases and phosphatases. *Front Biosci.* 2001 Oct 1;6:D1358-68. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11578967>
- Shabo G, Scheffer H, Cruysberg JR, Lammens M, Pasman JW, Spruit M, Willemsen MA. Congenital cataract facial dysmorphism neuropathy syndrome: a clinically recognizable entity. *Pediatr Neurol.* 2005 Oct;33(4):277-9.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16194727>
- Varon R, Gooding R, Steglich C, Marns L, Tang H, Angelicheva D, Yong KK, Ambrugger P, Reinhold A, Morar B, Baas F, Kwa M, Tournev I, Guerguelcheva V, Kremensky I, Lochmüller H, Müllner-Eidenböck A, Merlini L, Neumann L, Bürger J, Walter M, Swoboda K, Thomas PK, von Moers A, Risch N, Kalaydjieva L. Partial deficiency of the C-terminal-domain phosphatase of RNA polymerase II is associated with congenital cataracts facial dysmorphism neuropathy syndrome. *Nat Genet.* 2003 Oct;35(2):185-9. Epub 2003 Sep 21.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14517542>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/CTDP1>

Reviewed: April 2010
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services